


Applying the patch



Step 1

Select the area where OXYTROL® will be applied: abdomen, hip or buttock. Make sure skin is clean and dry.




Step 2

Carefully tear open the pouch and remove the OXYTROL® patch.



Step 3

Turn over and peel off half of the protective liner. Try not to touch the adhesive.



Step 4

Press the patch firmly on the skin, adhesive side down.



Step 5

Carefully remove the remaining half of the protective liner.



Step 6

Smooth the patch over your skin and press firmly. Wash your hands thoroughly.



The OXYTROL® patch should be changed on the same two days every week (eg. every Monday and Thursday) to maintain its beneficial effect.

Removing the patch

Remove old patches slowly and carefully to avoid skin damage. Once off, fold the patch in half for disposal. Warm water and mild soap can be used to remove the 'ring' of adhesive that may remain on the skin. A small amount of baby oil may also be used to remove any excess adhesive. Alcohol or similar liquids may cause skin irritation and should not be used. For further information about your condition or treatment please consult your healthcare professional.

Actavis 117 Harrington Street The Rocks Sydney 2000. Ph: 1300 881 893



Additional Information

To help keep the site of application healthy and irritation free;

- You can apply baby oil to the area, allow to dry and then apply the patch
- Rotate the site of application
- Remove any residual gum when removing patch
- If irritation and redness, apply 'over the counter' hydrocortisone cream to the area regularly for 5 days

Malvern Hill Consulting Suites - 1039 Malvern Road, Toorak, 3142

P: 03 9188 3690 F: 03 8513 6292